

```

=> file reg
=> s WRWWW/sqsp
L1      497 WRWWW/SQSP

=> s l1 and 4<=sql<=15
      1678496 4<=SQL<=15
L2      24 L1 AND 4<=SQL<=15

=> s WRWWW/sqsp and 4<=sql<=15
      24 WRWWW/SQSP
      1678496 4<=SQL<=15
L3      24 WRWWW/SQSP AND 4<=SQL<=15

=> file hcaplus

=> s l1 and pd<20040204
      19 L1
      24813213 PD<20040204
      (PD<20040204)
L4      6 L1 AND PD<20040204

=> dup rem l4
PROCESSING COMPLETED FOR L4
L5      3 DUP REM L4 (3 DUPLICATES REMOVED)

=> d l5 1-3 bib ab

L5      ANSWER 1 OF 3 HCAPLUS COPYRIGHT 2009 ACS on STN
AN      2004:502191 HCAPLUS <<LOGINID::20090202>>
DN      141:122298
TI      Identification of Peptides That Antagonize Formyl Peptide Receptor-Like
AU      1-Mediated Signaling
AU      Bae, Yoe-Sik; Lee, Ha Young; Jo, Eun Jin; Kim, Jung Im; Kang, Hyun-Kyu;
CS      Ye, Richard D.; Kwak, Jong-Young; Ryu, Sung Ho
CS      Medical Research Center for Cancer Molecular Therapy and Department of
        Biochemistry, College of Medicine, Dong-A University, Pusan, 602-714, S.
        Korea
SO      Journal of Immunology ( ***2004*** ), 173(1), 607-614
        CODEN: JOIMA3; ISSN: 0022-1767
PB      American Association of Immunologists
DT      Journal
LA      English
AB      Formyl peptide receptor-like 1 (FPR1) is an important classical
        chemoattractant receptor that is expressed in phagocytic cells in the
        peripheral blood and brain. Recently, various novel agonists have been
        identified from several origins, such as host-derived mols. Activation of
        FPR1 is closely related to inflammatory responses in the host defense
        mechanism and neurodegenerative disorders. Here, the authors identified
        several novel peptides by screening hexapeptide libraries that inhibit the
        binding of one of FPR1 agonists [Trp-Lys-Tyr-Met-Val-D-Met-CONH2
        (WKYMVM)] to its specific receptor, FPR1, in RBL-2H3 cells. Among the
        novel peptides, Trp-Arg-Trp-Trp-Trp-CONH2 [WRWWW (WRW4)] showed the
        most potent activity in terms of inhibiting WKYMVM binding to FPR1. The
        authors also found that WRW4 inhibited the activation of FPR1 by WKYMVM,
        resulting in the complete inhibition of the intracellular calcium
        increase, extracellular signal-regulated kinase activation, and
        chemotactic migration of cells toward WKYMVM. For the receptor
        specificity of WRW4 to the FPR family, the authors obsd. that WRW4
        specifically inhibit the increase in intracellular calcium by the FPR1
        agonists MMK-1, amyloid .beta.42 (A.beta.42) peptide, and F peptide, but
        not by the FPR agonist, fMLF. To investigate the effect of WRW4 on
        endogenous FPR1 ligand-induced cellular responses, the authors examd. its
        effect on A.beta.42 peptide in human neutrophils. A.beta.42
        peptide-induced superoxide generation and chemotactic migration of
        neutrophils were inhibited by WRW4, which also completely inhibited the
        internalization of A.beta.42 peptide in human macrophages. WRW4 is the
        first specific FPR1 antagonist and is expected to be useful in the study
        of FPR1 signaling and in the development of drugs against FPR1-related
        diseases.
RE.CNT  41      THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD
        ALL CITATIONS AVAILABLE IN THE RE FORMAT

L5      ANSWER 2 OF 3 HCAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 1
AN      2000:754713 HCAPLUS <<LOGINID::20090202>>
DN      133:330539

```

Sequence-determined DNA fragments and corresponding encoded polypeptides  
 from corn and Arabidopsis  
 Alexandrov, Nikolai; Brover, Vyacheslav; Chen, Xianfeng; Subramanian,  
 Gopalakrishnan; Troukhan, Maxim E.; Zheng, Liansheng; Dumas, J.  
 Ceres Inc., USA  
 Eur. Pat. Appl., 339 pp.  
 CODEN: EPXXDW  
 Patent  
 English  
 FAN.CNT 43

	PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
PI	EP 1033405	A2	20000906	EP 2000-301439	20000225 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	CA 2300692	A1	20000825	CA 2000-2300692	20000225 <--
	CA 2302828	A1	20001006	CA 2000-2302828	20000406 <--
	EP 1055728	A2	20001129	EP 2000-303770	20000504 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
	EP 1054060	A2	20001122	EP 2000-304161	20000517 <--
	R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
PRAI	US 1999-121825P	P	19990225		
	US 1999-145918P	P	19990727		
	US 1999-145951P	P	19990728		
	US 1999-146386P	P	19990802		
	US 1999-146388P	P	19990802		
	US 1999-146389P	P	19990802		
	US 1999-147038P	P	19990803		
	US 1999-147204P	P	19990804		
	US 1999-147302P	P	19990804		
	US 1999-147192P	P	19990805		
	US 1999-147260P	P	19990805		
	US 1999-147303P	P	19990806		
	US 1999-147416P	P	19990806		
	US 1999-147493P	P	19990809		
	US 1999-147935P	P	19990809		
	US 1999-148171P	P	19990810		
	US 1999-148319P	P	19990811		
	US 1999-148341P	P	19990812		
	US 1999-148565P	P	19990813		
	US 1999-148684P	P	19990813		
	US 1999-123180P	P	19990305		
	US 1999-123548P	P	19990309		
	US 1999-125788P	P	19990323		
	US 1999-126264P	P	19990325		
	US 1999-126785P	P	19990329		
	US 1999-127462P	P	19990401		
	US 1999-128234P	P	19990406		
	US 1999-128714P	P	19990408		
	US 1999-129845P	P	19990416		
	US 1999-130077P	P	19990419		
	US 1999-130449P	P	19990421		
	US 1999-130510P	P	19990423		
	US 1999-130891P	P	19990423		
	US 1999-131449P	P	19990428		
	US 1999-132048P	P	19990430		
	US 1999-132407P	P	19990430		
	US 1999-132484P	P	19990504		
	US 1999-132485P	P	19990505		
	US 1999-132486P	P	19990506		
	US 1999-132487P	P	19990506		
	US 1999-132863P	P	19990507		
	US 1999-134256P	P	19990511		
	US 1999-134218P	P	19990514		
	US 1999-134219P	P	19990514		
	US 1999-134221P	P	19990514		
	US 1999-134370P	P	19990514		
	US 1999-134768P	P	19990518		
	US 1999-134941P	P	19990519		
	US 1999-135124P	P	19990520		
	US 1999-135353P	P	19990521		
	US 1999-135629P	P	19990524		
	US 1999-136021P	P	19990525		
	US 1999-136392P	P	19990527		
	US 1999-136782P	P	19990528		
	US 1999-137222P	P	19990601		
	US 1999-137528P	P	19990603		

US 1999-137502P P 19990604  
US 1999-137724P P 19990607  
US 1999-138094P P 19990608

AB The present invention provides DNA mols. that constitute fragments of the genome and cDNAs from *Zea mays* (HYBRID SEED #35A19) and *Arabidopsis thaliana* (ecotype Wassilewsky), and polypeptides encoded thereby. The DNA mols. are useful for specifying a gene product in cells, either as a promoter or as a protein coding sequence or as an UTR or as a 3' termination sequence, and are also useful in controlling the behavior of a gene in the chromosome, in controlling the expression of a gene or as tools for genetic mapping, recognizing or isolating identical or related DNA fragments, or identification of a particular individual organism, or for clustering of a group of organisms with a common trait. *Arabidopsis* DNA is used in the present expt., but the procedure is a general one. Protocols are provided for Southern hybridizations and transformation of carrot cells. [This abstr. record is one of 15 records supplemental to CA1316218528Q necessitated by the large no. of index entries required to fully index the document and publication system constraints.].

L5 ANSWER 3 OF 3 HCAPLUS COPYRIGHT 2009 ACS on STN  
AN 2000:9190 HCAPLUS <<LOGINID::20090202>>  
DN 132:103595  
TI Sequence and analysis of chromosome 4 of the plant *Arabidopsis thaliana*  
AU Mayer, K.; Schuller, C.; Wambutt, R.; Murphy, G.; Volckaert, G.; Pohl, T.; Dusterhoft, A.; Stiekema, W.; Entlan, K.-D.; Terryn, N.; Harris, B.; Anson, V.; Brandt, P.; Grivell, L.; Rieger, M.; Weichselgartner, M.; De Simone, W.; Obermaier, B.; Mache, R.; Muller, M.; Kreis, M.; Delsen, M.; Pulgden, P.; Watson, M.; Schmidheini, T.; Reichert, B.; Portet, D.; Perez-Alonso, M.; Boutry, M.; Bancroft, I.; Vos, P.; Hoheisel, J.; Zimmermann, W.; Wedler, H.; Ridley, P.; Langham, S.-A.; McCullagh, B.; Bilham, L.; Robben, J.; Van Der Schueren, J.; Grymonprez, B.; Chuang, Y.-J.; Vandenbussche, F.; Braeken, M.; Weltjens, I.; Voet, M.; Bastiaens, I.; Aert, R.; Defoor, E.; Weitzenecker, T.; Bothe, G.; Ramsperger, U.; Hilbert, H.; Braun, M.; Holzer, E.; Brandt, A.; Peters, S.; Van Staveren, M.; Dirkse, W.; Mooijman, P.; Klein Lankhorst, R.; Rose, M.; Haut, J.; Kotter, P.; Berner, S.; Hempel, S.; Feldpausch, M.; Lamberth, S.; Van Den Daele, H.; De Keyser, A.; Buysschaert, C.; Gielen, J.; Villarroel, R.; De Clercq, R.; Van Montagu, M.; Rogers, J.; Cronin, A.; Quail, M.; Bray-Allen, S.; Clark, L.; Doggett, J.; Hall, S.; Kay, M.; Lennard, N.; McLay, K.; Mayes, R.; Pettett, A.; Rajandream, M.-A.; Lyne, M.; Benes, V.; Rechmann, S.; Borkova, D.; Blocker, H.; Scharfe, M.; Grimm, M.; Lohner, T.-H.; Dose, S.; De Haan, M.; Maarse, A.; Schafer, M.; Muller-Auer, S.; Gabel, C.; Fuchs, M.; Fartmann, B.; Granderath, K.; Dauner, D.; Herzl, A.; Neumann, S.; Argiriou, A.; Vitale, D.; Liguori, R.; Piravandi, E.; Massenet, O.; Quigley, P.; Clabaud, G.; Mundlein, A.; Felber, R.; Schnabl, S.; Hiller, R.; Schmidt, W.; Lecharny, A.; Aubourg, S.; Chedfor, F.; Cooke, R.; Berger, C.; Montfort, M.; Casacuberta, E.; Gibbons, T.; Weber, N.; Vandenbol, M.; Bagues, M.; Terol, J.; Torres, A.; Perez-Perez, A.; Purnelle, B.; Bent, E.; Johnson, S.; Tacon, D.; Jesse, T.; Heijnen, L.; Schwarz, S.; Scholler, P.; Heber, S.; Francis, P.; Biele, C.; Frishman, D.; Haase, D.; Lemcke, K.; Mewes, H. W.; Stocker, S.; Zaccaria, P.; Bevan, M.; Wilson, R. K.; De La Bastide, M.; Habermann, K.; Parnell, L.; Dedhia, N.; Gnoj, L.; Schutz, K.; Huang, E.; Spiegel, L.; Sehnk, M.; Murray, J.; Sheet, P.; Cordes, M.; Abu-Threideh, J.; Stoneking, T.; Kalicki, J.; Graves, T.; Harmon, G.; Edwards, J.; Latrelle, P.; Courtney, L.; Cloud, J.; Abbott, A.; Scott, K.; Johnson, D.; Minx, P.; Bentley, D.; Fulton, B.; Miller, N.; Greco, T.; Kemp, K.; Kramer, J.; Fulton, L.; Mardis, E.; Dante, M.; Pepin, K.; Hillier, L.; Nelson, J.; Spieth, J.; Ryan, E.; Andrews, S.; Geisel, C.; Layman, D.; Du, H.; Ali, J.; Berghoff, A.; Jones, K.; Drone, K.; Cotton, N.; Joshi, C.; Antoniou, B.; Zidanic, M.; Strong, C.; Sun, H.; Lamar, B.; Yordan, C.; Ma, P.; Zhong, J.; Preston, R.; Vil, D.; Shekher, M.; Matero, A.; Shah, R.; Swaby, I.K.; O'Shaughnessy, A.; Rodriguez, M.; Hoffman, J.; Till, S.; Granat, S.; Shoddy, N.; Hasegawa, A.; Hameed, A.; Lodhi, M.; Johnson, A.; Chen, E.; Marra, N.; Martienssen, R.; McCombie, W. R.

CS GSF-Forschungszentrum f. Umwelt u. Gesundheit, Munich Information Center for Protein Sequences am Max-Planck-Institut f. Biochemie, D-82152, Germany

SO Nature (London) ( \*\*\*\*1999\*\*\*\* ), 402(6763), 769-777  
CODEN: NATUAS; ISSN: 0028-0836  
PB Macmillan Magazines  
DT Journal  
LA English  
AB The higher plant *Arabidopsis thaliana* is an important model for identifying plant genes and detg. their function. To assist biol. investigations and to define chromosome structure, a coordinated effort to sequence the *Arabidopsis* genome was initiated in late 1996. This report describes one of the first milestones of this project, the sequence of chromosome 4. Anal. of 17.38 megabases of unique sequence, representing

about 17% of the genome, reveals 3744 protein coding genes, 81 tRNAs, and numerous repeat elements. Heterochromatic regions surrounding the putative centromere, which has not yet been completely sequenced, are characterized by an increased frequency of a variety of repeats, new repeats, reduced recombination, lowered gene d., and lowered gene expression. Roughly 60% of the predicted protein-coding genes have been functionally characterized on the basis of their homol. to known genes. Many genes encode predicted proteins that are homologous to human and *Caenorhabditis elegans* proteins.

RE.CNT 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s 13 and pd<20040204

10 L3

24813213 PD<20040204

(PD<20040204)

L6 1 L3 AND PD<20040204

=> d 16 bib ab

L6 ANSWER 1 OF 1 HCAPLUS COPYRIGHT 2009 ACS on STN

AN 2004:502191 HCAPLUS <<LOGINID::20090202>>

DN 141:122298

TI Identification of Peptides That Antagonize Formyl Peptide Receptor-Like 1-Mediated Signaling

AU Bae, Yoe-Sik; Lee, Ha Young; Jo, Eun Jin; Kim, Jung Im; Kang, Hyun-Kyu; Ye, Richard D.; Kwak, Jong-Young; Ryu, Sung Ho

CS Medical Research Center for Cancer Molecular Therapy and Department of Biochemistry, College of Medicine, Dong-A University, Pusan, 602-714, S. Korea

SO Journal of Immunology ( \*\*\*2004\*\*\* ), 173(1), 607-614

CODEN: JOIMA3; ISSN: 0022-1767

PB American Association of Immunologists

DT Journal

LA English

AB Formyl peptide receptor-like 1 (FPR1) is an important classical chemoattractant receptor that is expressed in phagocytic cells in the peripheral blood and brain. Recently, various novel agonists have been identified from several origins, such as host-derived mols. Activation of FPR1 is closely related to inflammatory responses in the host defense mechanism and neurodegenerative disorders. Here, the authors identified several novel peptides by screening hexapeptide libraries that inhibit the binding of one of FPR1 agonists [Trp-Lys-Tyr-Met-Val-D-Met-CONH2 (WKYMVM)] to its specific receptor, FPR1, in RBL-2H3 cells. Among the novel peptides, Trp-Arg-Trp-Trp-Trp-CONH2 [WRWWWW (WRW4)] showed the most potent activity in terms of inhibiting WKYMVM binding to FPR1. The authors also found that WRW4 inhibited the activation of FPR1 by WKYMVM, resulting in the complete inhibition of the intracellular calcium increase, extracellular signal-regulated kinase activation, and chemotactic migration of cells toward WKYMVM. For the receptor specificity of WRW4 to the FPR family, the authors obsd. that WRW4 specifically inhibit the increase in intracellular calcium by the FPR1 agonists MMK-1, amyloid .beta.42 (A.beta.42) peptide, and F peptide, but not by the FPR agonist, fMLP. To investigate the effect of WRW4 on endogenous FPR1 ligand-induced cellular responses, the authors examd. its effect on A.beta.42 peptide in human neutrophils. A.beta.42 peptide-induced superoxide generation and chemotactic migration of neutrophils were inhibited by WRW4, which also completely inhibited the internalization of A.beta.42 peptide in human macrophages. WRW4 is the first specific FPR1 antagonist and is expected to be useful in the study of FPR1 signaling and in the development of drugs against FPR1-related diseases.

RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> file .biotech

FILE 'MEDLINE' ENTERED AT 13:30:07 ON 02 FEB 2009

FILE 'CAPLUS' ENTERED AT 13:30:07 ON 02 FEB 2009

USE IS SUBJECT TO THE TERMS OF YOUR STN CUSTOMER AGREEMENT.

PLEASE SEE "HELP USAGETERMS" FOR DETAILS.

COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'BIOSIS' ENTERED AT 13:30:07 ON 02 FEB 2009

Copyright (c) 2009 The Thomson Corporation

FILE 'BIOTECHDS' ENTERED AT 13:30:07 ON 02 FEB 2009

FILE 'EMBASE' ENTERED AT 13:30:07 ON 02 FEB 2009  
Copyright (c) 2009 Elsevier B.V. All rights reserved.

FILE 'USPATFULL' ENTERED AT 13:30:07 ON 02 FEB 2009  
CA INDEXING COPYRIGHT (C) 2009 AMERICAN CHEMICAL SOCIETY (ACS)

FILE 'WPIDS' ENTERED AT 13:30:07 ON 02 FEB 2009  
COPYRIGHT (C) 2009 THOMSON REUTERS

=> s l3 and pd<20040204  
1 FILES SEARCHED...  
'SQL' NOT A VALID FIELD CODE  
'SQSP' IS NOT A VALID FIELD CODE  
4 FILES SEARCHED...  
'SQL' NOT A VALID FIELD CODE  
'SQSP' IS NOT A VALID FIELD CODE  
L7 1 L3 AND PD<20040204

=> d l7 bib ab

L7 ANSWER 1 OF 1 CAPLUS COPYRIGHT 2009 ACS on STN  
AN 2004:502191 CAPLUS <<LOGINID:20090202>>  
DN 141:122298  
TI Identification of Peptides That Antagonize Formyl Peptide Receptor-Like  
1-Mediated Signaling  
AU Bae, Yoe-Sik; Lee, Ha Young; Jo, Eun Jin; Kim, Jung Im; Kang, Hyun-Kyu;  
Ye, Richard D.; Kwak, Jong-Young; Ryu, Sung Ho  
CS Medical Research Center for Cancer Molecular Therapy and Department of  
Biochemistry, College of Medicine, Dong-A University, Pusan, 602-714, S.  
Korea  
SO Journal of Immunology ( \*\*\*2004\*\*\* ), 173(1), 607-614  
CODEN: JOIMA3; ISSN: 0022-1767  
PB American Association of Immunologists  
DT Journal  
LA English  
AB Formyl peptide receptor-like 1 (FPR1) is an important classical  
chemoattractant receptor that is expressed in phagocytic cells in the  
peripheral blood and brain. Recently, various novel agonists have been  
identified from several origins, such as host-derived moles. Activation of  
FPR1 is closely related to inflammatory responses in the host defense  
mechanism and neurodegenerative disorders. Here, the authors identified  
several novel peptides by screening hexapeptide libraries that inhibit the  
binding of one of FPR1 agonists [Trp-Lys-Tyr-Met-Val-D-Met-CONH2  
(WKYMVM)] to its specific receptor, FPR1, in RBL-2H3 cells. Among the  
novel peptides, Trp-Arg-Trp-Trp-Trp-Trp-CONH2 [WRW4] showed the  
most potent activity in terms of inhibiting WKYMVM binding to FPR1. The  
authors also found that WRW4 inhibited the activation of FPR1 by WKYMVM,  
resulting in the complete inhibition of the intracellular calcium  
increase, extracellular signal-regulated kinase activation, and  
chemotactic migration of cells toward WKYMVM. For the receptor  
specificity of WRW4 to the FPR family, the authors obsd. that WRW4  
specifically inhibit the increase in intracellular calcium by the FPR1  
agonists MMK-1, amyloid .beta.42 (A.beta.42) peptide, and F peptide, but  
not by the FPR agonist, fMLF. To investigate the effect of WRW4 on  
endogenous FPR1 ligand-induced cellular responses, the authors examd. its  
effect on A.beta.42 peptide in human neutrophils. A.beta.42  
peptide-induced superoxide generation and chemotactic migration of  
neutrophils were inhibited by WRW4, which also completely inhibited the  
internalization of A.beta.42 peptide in human macrophages. WRW4 is the  
first specific FPR1 antagonist and is expected to be useful in the study  
of FPR1 signaling and in the development of drugs against FPR1-related  
diseases.

RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD  
ALL CITATIONS AVAILABLE IN THE RE FORMAT

=> s l1 and pd<20040204  
1 FILES SEARCHED...  
'SQSP' IS NOT A VALID FIELD CODE  
4 FILES SEARCHED...  
'SQSP' IS NOT A VALID FIELD CODE  
L8 6 L1 AND PD<20040204

=> dup rem l8  
PROCESSING COMPLETED FOR L8  
L9 3 DUP REM L8 (3 DUPLICATES REMOVED)

L9 ANSWER 1 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN  
 AN 2004:502191 CAPLUS <<LOGINID::20090202>>  
 DN 141:122298  
 TI Identification of Peptides That Antagonize Formyl Peptide Receptor-Like  
 1-Mediated Signaling  
 AU Bae, Yoe-Sik; Lee, Ha Young; Jo, Eun Jin; Kim, Jung Im; Kang, Hyun-Kyu;  
 CS Ye, Richard D.; Kwak, Jong-Young; Ryu, Sung Ho  
 Medical Research Center for Cancer Molecular Therapy and Department of  
 Biochemistry, College of Medicine, Dong-A University, Pusan, 602-714, S.  
 Korea  
 SO Journal of Immunology ( \*\*\*2004\*\*\* ), 173(1), 607-614  
 CODEN: JOIMA3; ISSN: 0022-1767  
 PB American Association of Immunologists  
 DT Journal  
 LA English  
 AB Formyl peptide receptor-like 1 (FPR1) is an important classical  
 chemoattractant receptor that is expressed in phagocytic cells in the  
 peripheral blood and brain. Recently, various novel agonists have been  
 identified from several origins, such as host-derived mols. Activation of  
 FPR1 is closely related to inflammatory responses in the host defense  
 mechanism and neurodegenerative disorders. Here, the authors identified  
 several novel peptides by screening hexapeptide libraries that inhibit the  
 binding of one of FPR1 agonists [Trp-Lys-Tyr-Met-Val-D-Met-CONH2  
 (WKYMVM)] to its specific receptor, FPR1, in RBL-2H3 cells. Among the  
 novel peptides, Trp-Arg-Trp-Trp-Trp-Trp-CONH2 [WRWWWW (WRW4)] showed the  
 most potent activity in terms of inhibiting WKYMVM binding to FPR1. The  
 authors also found that WRW4 inhibited the activation of FPR1 by WKYMVM,  
 resulting in the complete inhibition of the intracellular calcium  
 increase, extracellular signal-regulated kinase activation, and  
 chemotactic migration of cells toward WKYMVM. For the receptor  
 specificity of WRW4 to the FPR family, the authors obsd. that WRW4  
 specifically inhibit the increase in intracellular calcium by the FPR1  
 agonists MMK-1, amyloid .beta.42 (A.beta.42) peptide, and F peptide, but  
 not by the FPR agonist, fMLF. To investigate the effect of WRW4 on  
 endogenous FPR1 ligand-induced cellular responses, the authors examd. its  
 effect on A.beta.42 peptide in human neutrophils. A.beta.42  
 peptide-induced superoxide generation and chemotactic migration of  
 neutrophils were inhibited by WRW4, which also completely inhibited the  
 internalization of A.beta.42 peptide in human macrophages. WRW4 is the  
 first specific FPR1 antagonist and is expected to be useful in the study  
 of FPR1 signaling and in the development of drugs against FPR1-related  
 diseases.

RE.CNT 41 THERE ARE 41 CITED REFERENCES AVAILABLE FOR THIS RECORD  
 ALL CITATIONS AVAILABLE IN THE RE FORMAT

L9 ANSWER 2 OF 3 CAPLUS COPYRIGHT 2009 ACS on STN DUPLICATE 1  
 AN 2000:754713 CAPLUS <<LOGINID::20090202>>  
 DN 133:330539  
 TI Sequence-determined DNA fragments and corresponding encoded polypeptides  
 from corn and Arabidopsis  
 IN Alexandrov, Nikolai; Brover, Vyacheslav; Chen, Xianfeng; Subramanian,  
 PA Gopalakrishnan; Troukhan, Maxim E.; Zheng, Liansheng; Dumas, J.  
 SO Ceres Inc., USA  
 Eur. Pat. Appl., 339 pp.  
 CODEN: EPKXDW  
 DT Patent  
 LA English  
 FAN.CNT 43

PATENT NO.	KIND	DATE	APPLICATION NO.	DATE
EP 1033405	A2	20000906	EP 2000-301439	20000225 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
CA 2300692	A1	20000825	CA 2000-2300692	20000225 <--
CA 2302828	A1	20001006	CA 2000-2302828	20000406 <--
EP 1055728	A2	20001129	EP 2000-303770	20000504 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
EP 1054060	A2	20001122	EP 2000-304161	20000517 <--
R: AT, BE, CH, DE, DK, ES, FR, GB, GR, IT, LI, LU, NL, SE, MC, PT, IE, SI, LT, LV, FI, RO				
US 1999-121825P	P	19990225		
US 1999-145918P	P	19990727		
US 1999-145951P	P	19990728		
US 1999-146386P	P	19990802		
US 1999-146388P	P	19990802		
US 1999-146389P	P	19990802		

	US 1999-147038P	P	19990803
	US 1999-147204P	P	19990804
	US 1999-147302P	P	19990804
	US 1999-147192P	P	19990805
	US 1999-147260P	P	19990805
	US 1999-147303P	P	19990806
	US 1999-147416P	P	19990806
	US 1999-147493P	P	19990809
	US 1999-147935P	P	19990809
	US 1999-148171P	P	19990810
	US 1999-148319P	P	19990811
	US 1999-148341P	P	19990812
	US 1999-148565P	P	19990813
	US 1999-148684P	P	19990813
	US 1999-123180P	P	19990305
	US 1999-123548P	P	19990309
	US 1999-125788P	P	19990323
	US 1999-126264P	P	19990325
	US 1999-126785P	P	19990329
	US 1999-127462P	P	19990401
	US 1999-128234P	P	19990406
	US 1999-128714P	P	19990408
	US 1999-129845P	P	19990416
	US 1999-130077P	P	19990419
	US 1999-130449P	P	19990421
	US 1999-130510P	P	19990423
	US 1999-130891P	P	19990423
	US 1999-131449P	P	19990428
	US 1999-132048P	P	19990430
	US 1999-132407P	P	19990430
	US 1999-132484P	P	19990504
	US 1999-132485P	P	19990505
	US 1999-132486P	P	19990506
	US 1999-132487P	P	19990506
	US 1999-132863P	P	19990507
	US 1999-134256P	P	19990511
	US 1999-134218P	P	19990514
	US 1999-134219P	P	19990514
	US 1999-134221P	P	19990514
	US 1999-134370P	P	19990514
	US 1999-134768P	P	19990518
	US 1999-134941P	P	19990519
	US 1999-135124P	P	19990520
	US 1999-135353P	P	19990521
	US 1999-135629P	P	19990524
	US 1999-136021P	P	19990525
	US 1999-136392P	P	19990527
	US 1999-136782P	P	19990528
	US 1999-137222P	P	19990601
	US 1999-137528P	P	19990603
	US 1999-137502P	P	19990604
	US 1999-137724P	P	19990607
	US 1999-138094P	P	19990608

AB The present invention provides DNA mols. that constitute fragments of the genome and cDNAs from Zea mays (HYBRID SEED #35A19) and Arabidopsis thaliana (ecotype Wassilewsky), and polypeptides encoded thereby. The DNA mols. are useful for specifying a gene product in cells, either as a promoter or as a protein coding sequence or as an UTR or as a 3' termination sequence, and are also useful in controlling the behavior of a gene in the chromosome, in controlling the expression of a gene or as tools for genetic mapping, recognizing or isolating identical or related DNA fragments, or identification of a particular individual organism, or for clustering of a group of organisms with a common trait. Arabidopsis DNA is used in the present expt., but the procedure is a general one. Protocols are provided for Southern hybridizations and transformation of carrot cells. [This abstr. record is one of 15 records supplemental to CA13316218528Q necessitated by the large no. of index entries required to fully index the document and publication system constraints.].

L9 ANSWER 3 OF 3 CAPLUS COPYRIGHT 2009 ACS ON STN  
 AN 2000:9190 CAPLUS <<LOGINID:20090202>>  
 DN 132:103595

TI Sequence and analysis of chromosome 4 of the plant Arabidopsis thaliana  
 AU Mayer, K.; Schuller, C.; Wambutt, R.; Murphy, G.; Volckaert, G.; Pohl, T.; Dusterhoft, A.; Stiekema, W.; Entlan, K.-D.; Terryn, N.; Harris, B.; Ansonge, W.; Brandt, P.; Grivell, L.; Rieger, M.; Weichselgartner, M.; De Simone, V.; Obermaier, B.; Mache, R.; Muller, M.; Kreis, M.; Delseny, M.; Pulgden, P.; Watson, M.; Schmidheini, T.; Reichert, B.; Portetalle, D.; Perez-Alonso, M.; Boutry, M.; Bancroft, I.; Vos, P.; Hoheisel, J.;

Zimmermann, W.; Wedler, H.; Ridley, P.; Langham, S.-A.; McCullagh, B.; Bilham, L.; Robben, J.; Van Der Schueren, J.; Grymonprez, B.; Chuang, Y.-J.; Vandenbussche, F.; Braeken, M.; Weltjens, I.; Voet, M.; Bastiaens, I.; Aert, R.; Defoor, E.; Weitzenegger, T.; Bothe, G.; Ramsperger, U.; Hilbert, H.; Braun, M.; Holzer, E.; Brandt, A.; Peters, S.; Van Staveren, M.; Dirkse, W.; Mooijman, P.; Klein Lankhorst, R.; Rose, M.; Haut, J.; Kotter, P.; Berneiser, S.; Hempel, S.; Feldpausch, M.; Lamberth, S.; Van Den Daele, H.; De Keyser, A.; Buysshaert, C.; Gielen, J.; Villarroel, R.; De Clercq, R.; Van Montagu, M.; Rogers, J.; Cronin, A.; Quail, M.; Bray-Allen, S.; Clark, L.; Doggett, J.; Hall, S.; Kay, M.; Lennard, N.; McClay, K.; Mayes, R.; Pettett, A.; Rajandream, M.-A.; Lyne, M.; Benes, V.; Rechmann, S.; Borkova, D.; Blocker, H.; Scharfe, M.; Grimm, M.; Lohnert, T.-H.; Dose, S.; De Haan, M.; Maarse, A.; Schafer, M.; Muller-Auer, S.; Gabel, C.; Fuchs, M.; Fartmann, B.; Granderath, K.; Dauner, D.; Herzl, A.; Neumann, S.; Argiriou, A.; Vitale, D.; Liguori, R.; Piravandi, E.; Massenet, O.; Quigley, F.; Clabaud, G.; Mundlein, A.; Felber, R.; Schnabl, S.; Hiller, R.; Schmidt, W.; Lecharny, A.; Aubourg, S.; Chefedor, F.; Cooke, R.; Berger, C.; Montfort, M.; Casacuberta, E.; Gibbons, T.; Weber, N.; Vandenbol, M.; Bagues, M.; Terol, J.; Torres, A.; Perez-Perez, A.; Purnelle, B.; Bent, E.; Johnson, S.; Tacon, D.; Jesse, T.; Heijnen, L.; Schwarz, S.; Scholler, P.; Heber, S.; Francis, P.; Bielek, C.; Frishman, D.; Haase, D.; Lemcke, K.; Mewes, H. W.; Stocker, S.; Zaccaria, P.; Bevan, M.; Wilson, R. K.; De La Bastide, M.; Habermann, K.; Parnell, L.; Dedhia, N.; Gnoj, L.; Schutz, K.; Huang, B.; Spiegel, L.; Sehkou, M.; Murray, J.; Sheet, P.; Cordes, M.; Abu-Threideh, J.; Stoneking, T.; Kalicki, J.; Graves, T.; Harmon, G.; Edwards, J.; Latrelle, P.; Courtney, L.; Cloud, J.; Abbott, A.; Scott, K.; Johnson, D.; Minx, P.; Bentley, D.; Fulton, B.; Miller, N.; Greco, T.; Kemp, K.; Kramer, J.; Fulton, L.; Mardis, E.; Dante, M.; Pepin, K.; Hillier, L.; Nelson, J.; Spieth, J.; Ryan, E.; Andrews, S.; Geisel, C.; Layman, D.; Du, H.; Ali, J.; Berghoff, A.; Jones, K.; Drone, K.; Cotton, N.; Joshua, C.; Antoniou, B.; Zidanic, M.; Strong, C.; Sun, H.; Lamar, B.; Yordan, C.; Ma, P.; Zhong, J.; Preston, R.; Vil, D.; Shekher, M.; Matero, A.; Shah, R.; Swaby, I'K.; O'Shaughnessy, A.; Rodriguez, M.; Hoffman, J.; Till, S.; Granat, S.; Shohdy, N.; Hasegawa, A.; Hameed, A.; Lodhi, M.; Johnson, A.; Chen, E.; Marra, M.; Martienssen, R.; McCombie, W. R.

CS GSF-Forschungszentrum f. Umwelt u. Gesundheit, Munich Information Center  
for Protein Sequences am Max-Planck-Institut f. Biochemie, D-82152,  
Germany

SO Nature (London) ( \*\*\*\*1999\*\*\* ), 402(6763), 769-777

CODEN: NATUAS; ISSN: 0028-0836

FB Macmillan Magazines

DT Journal

LA English

AB The higher plant *Arabidopsis thaliana* is an important model for identifying plant genes and detg. their function. To assist bio. investigations and to define chromosome structure, a coordinated effort to sequence the *Arabidopsis* genome was initiated in late 1996. This report describes one of the first milestones of this project, the sequence of chromosome 4. Anal. of 17.38 megabases of unique sequence, representing about 17% of the genome, reveals 3744 protein coding genes, 81 tRNAs, and numerous repeat elements. Heterochromatic regions surrounding the putative centromere, which has not yet been completely sequenced, are characterized by an increased frequency of a variety of repeats, new repeats, reduced recombination, lowered gene d., and lowered gene expression. Roughly 60% of the predicted protein-coding genes have been functionally characterized on the basis of their homol. to known genes. Many genes encode predicted proteins that are homologous to human and *Caenorhabditis elegans* proteins.

RE.CNT 50 THERE ARE 50 CITED REFERENCES AVAILABLE FOR THIS RECORD

ALL CITATIONS AVAILABLE IN THE RE FORMAT

=>  
--Logging off of STN--

=>  
Executing the logoff script...

=> LOG Y

STN INTERNATIONAL LOGOFF AT 13:34:24 ON 02 FEB 2009